



## TAFLEN WYBODAETH - INFORMATION SHEET

### SWEETCORN



**Establishment:** Sweet corn, like maize, will grow well on light free draining soils. Soil pH should be between 6.0 and 7.5. Sow into warm soil (10°C or higher) at a density of 60,000 – 70,000 seeds/ha and at a depth of 2.5 -3.5cm. Super sweet varieties must be grown separately from traditional corn or maize to prevent cross pollination.

**Nutrient requirements:** Approximately 40-50kg/ha of nitrogen should be available in the seedbed during the early stages of growth. Apply maintenance levels of phosphate and potash.

**Weeds, diseases and pests:** Atrazine or clopyralid based herbicides can be used if needed. Pests and diseases are few but can be severe. Smut is rare but cannot be treated.

**Harvesting date and method:** Harvest by hand or with a machine any time from early August onwards. It is important to grade cobs for sale if machine harvesting.

**Uses:** Suitable for farm gate, box schemes or pick your own sale. Using a mixture of varieties can increase period of availability.

**Processing and the market:** Trim cobs as soon as harvested and store at 0°C. Cobs will remain in a marketable state for no more than two weeks. Demand for locally produced vegetables is increasing.

### INDIA-CORN

**Sefydliad:** Bydd india-corn, fel corn (maize), yn tyfu'n dda mewn pridd ysgafn sy'n draenio'n rhydd. Dylai pH y pridd fod rhwng 6.0 a 7.5. Plannwch mewn pridd cynnes (10°C neu uwch) mewn dwysedd o 60,000 - 70,000 had/ha ac ar ddyfnder o 2.5 - 3.5cm. Rhaid tyfu mathau melys ar wahân i'r corn traddodiadol er mwyn osgoi trawsbeilliad.

**Anghenion maethol:** Dylai fod tua 40-50kg/ha o nitrogen ar gael yn y gwely had yn ystod cyfnod cynnar y tyfiant. Taenwch lefelau cynhaliaeth o ffosffad a photasiwm.

**Chwyn, afiechydon a phlâu:** Gellir defnyddio chwynladdwyr gyda sylfaen Atrazine neu clopyralid os oes angen. Ychydig o blâu ac afiechydon a geir ond gallant fod yn llym. Mae penddu (smut) yn brin ond ni ellir ei drin.

**Dyddiad a dull cynaeafu:** Cynaeafu â llaw neu gyda pheiriant ar unrhyw adeg o ddechrau Awst ymlaen. Mae'n bwysig graddio'r tywysennau ar gyfer gwerthiant os byddant yn cael eu cynaeafu â pheiriant.

**Defnyddiau:** Addas ar gyfer gwerthiant giât fferm, cynlluniau bocs neu werthiant pigo eich hunain. Drwy ddefnyddio cymysgedd o fathau, gellir cynyddu'r cyfnod y maent ar gael.

**Prosesu a'r farchnad:** Dylech docio'r tywysennau cyn gynted ag y byddwch yn eu cynaeafu a'u storio ar dymheredd o 0°C. Bydd y tywysennau'n cadw mewn cyflwr gwerthadwy am bythefnos yn unig. Mae'r galw am lysiau sydd wedi'u tyfu'n lleol yn cynyddu.

*Whilst every effort is made to ensure the information provided on this sheet is correct, CALU cannot be held responsible of any actions taken on the basis of its content.*

Tra gwneir pob ymdrech i sicrhau bod y wybodaeth yn y daflen hon yn gywir, ni ellir dal CALU'n gyfrifol am ganlyniadau unrhyw weithred a gymerir ar sail ei gynnwys.

CALU - ☎01248 680450 🌐 [www.calu.bangor.ac.uk](http://www.calu.bangor.ac.uk) ✉ [calu@bangor.ac.uk](mailto:calu@bangor.ac.uk)